
URCLIFT BPH Relief. In Sight...

Before and After



- Minimally invasive
- Minimal downtime¹
- Rapid symptom relief^{1,2}
- ntime¹ Preserves sexual function³
- Does not preclude other treatment options

1. Shore, Can J Urology 2014 2. Roehrborn, J Urology 2013 LIFT Study 3. McVary, J Sexual Medicine 2014 4. Chin, Urology 2012

www.UroLiftForBPH.com



UroLift[®] System Benefits Comparison

| Symptom Relief | UroLift®* | TURP | Laser | Thermotherapy | Medications |
|--|-----------|------|-------|---------------|-------------|
| Minimally invasive treatment | ~ | | | ~ | ~ |
| Rapid relief without complications ^{1,2} | ~ | | | | |
| Durable results after one treatment ^{2,4} | ✓ | ✓ | ✓ | ~ | |

Complications

| No cutting, heating, or removal of tissue | ~ | | | ✓ |
|--|---|--|--------------|---|
| No complications associated with major surgery/general anesthesia ² | ~ | | \checkmark | ✓ |
| Does not cause bleeding requiring transfusion ² | ~ | | ✓ | ✓ |
| Does not cause erectile dysfunction or ejaculatory dysfunction ^{2,3} | ~ | | | |
| Does not cause permanent urinary incontinence ² | ✓ | | ✓ | ✓ |

Convenience

| Doctor can perform procedure in the office/clinic (not OR) | \checkmark | | | ✓ | ✓ |
|--|--------------|---|---|---|---|
| Typically no catheter required after procedure | ✓ | | | | ✓ |
| No medications required to sustain relief ² | ✓ | ~ | ~ | ✓ | |

*Most common adverse events reported include hematuria, dysuria, micturition urgency, pelvic pain, and urge incontinence. Most symptoms were mild to moderate in severity and resolved within two to four weeks after the procedure.